

Assessing the Effects of Instrument Systematic Uncertainty in the IR Measurements on Derivation of Spectral Fingerprints Temperatures

Nipa Phojanamongkolkij, Marty Mlynczak, Dave Johnson, Seiji Kato,

Dave Kratz, Xu Liu, and Fred Rose

NASA Langley Research Center

CLARREO SDT Meeting

October 12-14th, 2011

Agenda

- Background/Objectives
- Detail Analysis
 - Input definition
 - Method
 - Output
- Conclusion
- Next Steps

Background


- ❖ Continuous work on determining an allowable IR systematic uncertainty distribution across the 200-2000 (cm^{-1}) wavenumbers and scene temperatures of 200-300K that will still enable the derivation of physical parameters.
- ❖ Unrealistic IR systematic uncertainty was used in the previous work.

Objectives

- ❖ To assess the effects of the realistic IR systematic uncertainty on the derivation of spectral fingerprints.
- ❖ Ultimately, to help in a what-if study on the IR uncertainty requirement effects on the fingerprint derivation.


Current Study

- Inputs
 - Datasets: the radiance differences of all-sky CCCMA from Huang et al. [2010].
 - Instrument bias function: IR instrument systematic uncertainty model from Dave Johnson by using scene temperatures from Seiji's TB zonal annual averages.
 - Baseline (no bias) instrument
 - To study a 10-year expected change, the radiance difference is decreased by a factor of 10.
 - CLARREO instrument
 - Ensembles of IR bias were simulated and added to the baseline data.
- Method - Huang et al. [2010]
- Outputs - Performance measures of % detection.



Three detectors were selected:


- a pyroelectric detector (Pyro) to cover 200-700 cm⁻¹
- a longwave (LW) HgCdTe detector (CrMCT) to cover 600-1400 cm⁻¹
- a midwave (MW) HgCdTe detector (MwMCT) to cover 1000-2000 cm⁻¹.




Three detectors were selected:

- a pyroelectric detector (Pyro) to cover 200-700 cm^{-1}
- a longwave (LW) HgCdTe detector (CrMCT) to cover 600-1400 cm^{-1}
- a midwave (MW) HgCdTe detector (MwMCT) to cover 1000-2000 cm^{-1} .


Current Instrument Bias ($k=1$)



Current Instrument Bias (k=1)




Signal Signatures – CCCMA model from Huang et al. [2010].




Fingerprint Example: All effects of Zone 10

(Blue = No Bias, Red = CLARREO)


Input $y = S_1 a_1 + S_2 a_2 + \dots + S_9 a_9 + \varepsilon$




Output $\hat{y} = S_1 \hat{a}_1 + S_2 \hat{a}_2 + \dots + S_9 \hat{a}_9$



Fitting Error (FE) = Input - Output = ε



Retrieval Uncertainty (RU) = $U(\hat{y})$



Fingerprint Uncertainty

Fitting Error

- Potential causes:
 - Missing nonlinearity of atmospheric effects.
 - Missing vertical resolution of atmospheric effects.
 - Missing atmospheric properties (e.g., cloud optical thickness, cloud height, cloud fraction, etc.)
- ? Other unknown

Retrieval Uncertainty


- Potential causes:
 - + Highly spectral, spatial, and temporal correlated data.
 - + High natural variability.
 - + High signal shape uncertainty.
 - + Missing atmospheric effects in the analysis.
- ? Other unknown

How ensembles of bias function were obtained?

Let N = Number of Simulation

$RN_{N \times 6} = \text{randn}(0,1)$

For $i = 1$ to N




End

where $\% \text{ Detection} = \% \text{ of Number of Signal} \geq \text{Threshold}$


Determining IR Bias Function

For simulation run i^{th} ,


Period 1:



Period 2:




Difference:




How many simulations (N)?

Simulation Error vs. Number of Simulations For All Zones



Perturbed all effects – No Bias Baseline


(Blue = Retrieval Mean of No Bias Baseline, Green = Fingerprint Uncertainty)



Define: $\text{Signal(No Bias)} = \text{Retrieval Mean of No Bias} / \text{Fingerprint Uncertainty of No Bias}$

Perturbed all effects - CLARREO

(Red = Retrieval Mean of CLARREO, Green = Fingerprint Uncertainty)




Define: $\text{Signal}(\text{CLARREO}) = \text{Retrieval Mean of CLARREO} / \text{Fingerprint Uncertainty of CLARREO}$

Perturbed all effects – No Bias Baseline vs. CLARREO

17


(Blue = Signal of No Bias, Red = Signal of CLARREO)



Define: % Detection = Relative volume of signals that are at least the specified threshold


% Detection Comparison

Blue = No Bias; Gray = Ensembles of CLARREO



%Detection by Effects (all zones combined)

Blue = No Bias; Gray = Ensembles of CLARREO



Conclusion/Next Steps

We have demonstrated a framework of applying fingerprinting (based on Huang et al. [2010]) to assess the effects of the realistic IR systematic uncertainty on the derivation of spectral fingerprints.

Next Steps

- Counter-intuitive on Retrieval Uncertainty (RU) is much smaller than the Fitting Error (FE).
 - Are we underestimating the RU?
 - Are we under-fitting (left out) atmospheric effects?
 - Or both?
- Try with a comprehensive and continuous climate dataset (multiple years, multiple vertical resolutions, etc.)